

THE CHEMISTRY INSIDE SPICES & HERBS: RESEARCH AND DEVELOPMENT

VOLUME 4



Editors:
Pankaj Kumar Chaurasia
Shashi Lata Bharati
Sunita Singh

Bentham Books

The Chemistry Inside Spices & Herbs: Research and Development

(Volume 4)

Edited by

Pankaj Kumar Chaurasia

*P.G. Department of Chemistry
L.S. College, B.R. Ambedkar Bihar University
Muzaffarpur-842001, Bihar, India*

Shashi Lata Bharati

*Department of Chemistry
North Eastern Regional Institute of Science and Technology
(NERIST) Nirjuli-791109
Arunachal Pradesh, India*

&

Sunita Singh

*Department of Chemistry
Navyug Kanya Mahavidyalaya, University of Lucknow
Lucknow- 226004
Uttar Pradesh, India*

The Chemistry Inside Spices & Herbs: Research and Development (Volume 4)

Editors: Pankaj Kumar Chaurasia, Shashi Lata Bharati and Sunita Singh

ISSN (Online): 5263/2; 54

ISSN (Print): 5263/2; 89

ISBN (Online): 978-981-5196-83-2

ISBN (Print): 978-981-5196-84-9

ISBN (Paperback): 978-981-5196-85-6

© 2024, Bentham Books imprint.

Published by Bentham Science Publishers Pte. Ltd. Singapore. All Rights Reserved.

First published in 2024.

BENTHAM SCIENCE PUBLISHERS LTD.

End User License Agreement (for non-institutional, personal use)

This is an agreement between you and Bentham Science Publishers Ltd. Please read this License Agreement carefully before using the book/echapter/ejournal (“**Work**”). Your use of the Work constitutes your agreement to the terms and conditions set forth in this License Agreement. If you do not agree to these terms and conditions then you should not use the Work.

Bentham Science Publishers agrees to grant you a non-exclusive, non-transferable limited license to use the Work subject to and in accordance with the following terms and conditions. This License Agreement is for non-library, personal use only. For a library / institutional / multi user license in respect of the Work, please contact: permission@benthamscience.net.

Usage Rules:

1. All rights reserved: The Work is the subject of copyright and Bentham Science Publishers either owns the Work (and the copyright in it) or is licensed to distribute the Work. You shall not copy, reproduce, modify, remove, delete, augment, add to, publish, transmit, sell, resell, create derivative works from, or in any way exploit the Work or make the Work available for others to do any of the same, in any form or by any means, in whole or in part, in each case without the prior written permission of Bentham Science Publishers, unless stated otherwise in this License Agreement.
2. You may download a copy of the Work on one occasion to one personal computer (including tablet, laptop, desktop, or other such devices). You may make one back-up copy of the Work to avoid losing it.
3. The unauthorised use or distribution of copyrighted or other proprietary content is illegal and could subject you to liability for substantial money damages. You will be liable for any damage resulting from your misuse of the Work or any violation of this License Agreement, including any infringement by you of copyrights or proprietary rights.

Disclaimer:

Bentham Science Publishers does not guarantee that the information in the Work is error-free, or warrant that it will meet your requirements or that access to the Work will be uninterrupted or error-free. The Work is provided "as is" without warranty of any kind, either express or implied or statutory, including, without limitation, implied warranties of merchantability and fitness for a particular purpose. The entire risk as to the results and performance of the Work is assumed by you. No responsibility is assumed by Bentham Science Publishers, its staff, editors and/or authors for any injury and/or damage to persons or property as a matter of products liability, negligence or otherwise, or from any use or operation of any methods, products instruction, advertisements or ideas contained in the Work.

Limitation of Liability:

In no event will Bentham Science Publishers, its staff, editors and/or authors, be liable for any damages, including, without limitation, special, incidental and/or consequential damages and/or damages for lost data and/or profits arising out of (whether directly or indirectly) the use or inability to use the Work. The entire liability of Bentham Science Publishers shall be limited to the amount actually paid by you for the Work.

General:

1. Any dispute or claim arising out of or in connection with this License Agreement or the Work (including non-contractual disputes or claims) will be governed by and construed in accordance with the laws of Singapore. Each party agrees that the courts of the state of Singapore shall have exclusive jurisdiction to settle any dispute or claim arising out of or in connection with this License Agreement or the Work (including non-contractual disputes or claims).
2. Your rights under this License Agreement will automatically terminate without notice and without the

need for a court order if at any point you breach any terms of this License Agreement. In no event will any delay or failure by Bentham Science Publishers in enforcing your compliance with this License Agreement constitute a waiver of any of its rights.

3. You acknowledge that you have read this License Agreement, and agree to be bound by its terms and conditions. To the extent that any other terms and conditions presented on any website of Bentham Science Publishers conflict with, or are inconsistent with, the terms and conditions set out in this License Agreement, you acknowledge that the terms and conditions set out in this License Agreement shall prevail.

Bentham Science Publishers Pte. Ltd.

80 Robinson Road #02-00

Singapore 068898

Singapore

Email: subscriptions@benthamscience.net



CONTENTS

PREFACE	i
LIST OF CONTRIBUTORS	iii
CHAPTER 1 ROLE OF INDIAN SPICES IN THE MANAGEMENT OF EMERGING VIRAL INFECTIONS	1
<i>Anuradha Singh</i>	
INTRODUCTION	1
Viral Life Cycle and Possible Drug Interventions	3
Traditional Plant-Based Remedies	5
Antiviral Potential of Indian Spices	6
Turmeric	7
Ginger (<i>Zingiber officinale</i> Roscoe)	9
Garlic (<i>Allium sativum</i>)	10
Black Pepper (<i>Piper nigrum</i>)	11
CONCLUSION	12
ACKNOWLEDGMENT	13
REFERENCES	13
CHAPTER 2 CHEMICAL, FUNCTIONAL, AND NUTRITIONAL PROPERTIES OF CORIANDER (<i>CORIANDRUM SATIVUM</i>)	19
<i>Rafael A. Zambelli</i>	
INTRODUCTION	19
Chemical Properties	21
Functional and Nutritional Properties	29
Applications	32
CONCLUSION	35
REFERENCES	36
CHAPTER 3 PHYTOCHEMICAL AND PHARMACOLOGICAL POTENTIAL OF <i>SPHAERANTHUS INDICUS</i> LINN	44
<i>Vaishali Tiwari, Varsha Yadav and Arjun Patra</i>	
INTRODUCTION	44
PHYTOCHEMISTRY AND BIOACTIVE COMPOUNDS OF <i>SPHAERANTHUS INDICUS</i>	46
PHARMACOLOGICAL ACTIVITIES OF <i>SPHAERANTHUS INDICUS</i>	51
Antioxidant Activity	51
Wound Healing Activity	53
Anxiolytic Activity	54
Antimicrobial Activity	54
Mast Cell Stabilizing Activity	59
Anti-cancer Activity	60
Antidiabetic Activity	60
Antihyperlipidemic Activity	62
Anti-inflammatory Activity	62
Bronchodilatory Activity	64
Larvicidal Activity	64
Analgesic Activity	65
Hepatoprotective Activity	66
Neuropharmacological Activity	68
Antiasthmatic Activity	69

Immunomodulatory Activity	69
Anticonvulsant Activity	70
Anti-viral Activity	70
Diuretic and Antiulcer Activity	70
Nephroprotective Activity	70
Antilithiatric Activity	71
Anti-amnesic Activity	71
Toxicity Studies	71
CONCLUSION	72
REFERENCES	72
CHAPTER 4 THE CHEMISTRY AND PHARMACOLOGICAL STUDY OF <i>COPAIFERA</i>	
SPP. OILS	79
<i>Rajesh Kumar, Yogendra Singh, Mona Piplani, Pankaj Bhateja, Ruchika Garg and Brijesh Kumar</i>	
INTRODUCTION	80
ECOLOGICAL DISTRIBUTION OF NOTABLE SPECIES	82
ETHNOPHARMACOLOGICAL USES	83
BIOACTIVE COMPOUNDS	85
MAJOR SESQUITERPENES FROM <i>COPAIFERA</i> OLEORESIN	93
MAJOR DITERPENES FROM <i>COPAIFERA</i> OLEORESIN	97
PHARMACOLOGICAL ACTIVITIES	99
Anti-inflammatory Activity	100
Wound Healing Activity	100
Antimicrobial Activity	101
Hepatoprotective Activity	102
Anticancer Activity	102
Antinociceptive Activity	102
Mutagenic and Cytotoxic Activity	102
Insecticidal Activity	103
Larvicidal Activity	103
Antileishmanial Activity	103
Miscellaneous Activities	104
ANALYTICAL STUDIES	104
SAFETY ASPECTS	106
CONCLUSION	108
REFERENCES	108
CHAPTER 5 PHARMACOGNOSTIC PROFILE OF A POTENT MEDICINAL HERB:	
<i>NARDOSTACHYS JATAMANSI</i>	117
<i>Palak Gupta, Neetika Naudiyal, Pratima Raypa and Vandana A. Kumar</i>	
INTRODUCTION	118
Botanical Classification	119
History of Use	120
Use in Ayurveda and Siddha	120
Use in Unani	121
Common Name	121
Morphology	121
Occurrence, Distribution and Cultivation	121
Endangered Status	122
Phytochemistry	122
Tissue Culture	124

Antioxidant Potential	126
Metabolic Profiling	128
Biological Importance of Nutrient Elements	130
Antimicrobial Activity	132
<i>In-vitro</i> Cytotoxic Potential	133
Anti-diabetic Potential	135
Anti-inflammatory Potential	136
CONCLUDING REMARK	137
ACKNOWLEDGEMENT	138
REFERENCES	138
CHAPTER 6 POTENTIALS OF WILD EDIBLE BUSH ONION (<i>AFROSTYRAX</i> <i>LEPIDOPHYLLUS</i>) PLANT AS A SOURCE OF ESSENTIAL FOOD AND MEDICINAL CONSTITUENTS	147
<i>Victor Olusegun Oyetayo, Folake Lucy Oyetayo and Nathaniel Olawale Oluji</i>	
INTRODUCTION	147
Botanical Description and Historical Cultivation	149
Chemical and Nutritional Composition of Bush Onion	149
Anti-nutrient Composition of the Bush Onion Plant	151
Food and Medicinal Uses of the Bush Onion Plant	151
Antimicrobial Activities of the Bush Onion Plant	152
Antioxidant Activities of the Bush Onion Plant	152
Anti-Cancer Activities of the Bush Onion Plant	153
Other Uses of the Bush Onion Plant	153
CONCLUSION AND RECOMMENDATIONS	153
REFERENCES	153
CHAPTER 7 ESSENTIAL OIL: ITS CHEMISTRY, BIOCHEMISTRY AND APPLICATION	157
<i>Mohit Agrawal, Komal Rao, Manmohan Singhal and Hema Chaudhary</i>	
INTRODUCTION	157
CHEMISTRY OF ESSENTIAL OILS (EOs)	158
Terpenoids	159
Oxygenated Compounds	160
Monoterpene / Monoterpenoid	160
Ketones	160
Esters	160
Extraction Methods of Essential Oils	161
<i>Cold-pressed Method</i>	161
<i>Steam-distillation Method</i>	161
<i>Solvent-extraction Method</i>	161
Mechanism Action of Essential Oils	161
Applications of Essential Oils	162
Lavender Oil	163
<i>Composition</i>	163
<i>Applications</i>	163
Clove Oil	164
<i>Applications</i>	164
Eucalyptus Oil	164
<i>Composition</i>	164
<i>Applications</i>	164
Peppermint Oil	165
<i>Properties</i>	165

Lemongrass Oil	165
Cinnamon Oil	165
Essential Oils in Aromatherapy	166
<i>Advantages of Aromatherapy</i>	166
<i>Essential Oils used in Aromatherapy</i>	167
<i>Side Effects Caused by Essential Oils</i>	168
CONCLUSION	168
REFERENCES	169
CHAPTER 8 THE PLANT PROTEASES-A GROUP OF POTENTIAL PHARMACOLOGICAL, BIOTECHNOLOGICAL, AND INDUSTRIALLY IMPORTANT ENZYMES	175
<i>Tina Roy, Sweta Arora, Shreyansi Das, Nilasish Pal and Nirmalendu Das</i>	
INTRODUCTION	176
CLASSIFICATION OF PROTEASES	177
Exopeptidases	178
<i>Aminopeptidases</i>	178
<i>Carboxypeptidases</i>	178
Endopeptidases	178
<i>Serine Proteases</i>	178
<i>Cysteine/Thiol Proteases</i>	179
<i>Metalloproteases</i>	179
<i>Aspartic Proteases</i>	179
<i>Threonine Proteases</i>	180
<i>Glutamic Proteases</i>	180
SOURCES OF PROTEASES	180
Plants	180
Animals	184
Microbes	184
CLASSIFICATION BY CATALYTIC TYPE	185
Catalytic Mechanism of Serine Protease	186
Catalytic Mechanism of Threonine Protease	187
Catalytic Mechanism of Cysteine Protease	188
Catalytic Mechanism of Aspartic Protease	189
Catalytic Mechanism of Glutamic Protease	190
Catalytic Mechanism of Metallo-protease	191
Self-cleaving Proteins	192
APPLICATION OF PROTEASES	193
Therapeutic	194
<i>Intestinal Disorders</i>	194
<i>Diabetic Thrombosis</i>	194
<i>Rheumatoid Arthritis</i>	194
<i>Wound Healing</i>	194
<i>Anticancer and Antitumor Activity</i>	195
<i>Other</i>	195
Industrial	196
<i>Meat Tenderization</i>	196
<i>Dairy Industry</i>	196
<i>Baking Industry</i>	197
<i>Animal Feed</i>	197
<i>Brewing and Wine Industry</i>	197

<i>Bioethanol Production</i>	197
<i>Cosmetics</i>	197
EXTRACTION	198
<i>IN VIVO</i> PROTEASE PRODUCTION	198
<i>IN VITRO</i> PROTEASE PRODUCTION	198
Micropropagation and Somatic Embryogenesis	200
Callus and Cell Suspension Culture	200
CONCLUSION	200
ACKNOWLEDGEMENT	201
REFERENCES	201
SUBJECT INDEX	234

PREFACE

VOLUME-4 of the book series “**The Chemistry Inside Spices and Herbs: Research and Development**” is a nice collection of a total of eight chapters written by experts from India, Brazil, and Nigeria. Volume-4 has chapters full of scientific knowledge on the pharmacology of different herbs and spices, and different pharmaceutical roles of herbs and spices along with their chemistry have been excellently expounded by the experts. This volume of the book includes several pharmaceutically relevant topics such as the pharmacological potential of *Sphaeranthus indicus* Linn, Indian spices in the management of emerging viral infections, Coriander and its properties, *Copaifera* spp. Oils’ pharmacological study, *Nardostachys jatamansi* and its pharmacognostic profile, edible bush onion (*Afrotyrax lepidophyllus*) as a source of essential food and medicinal constituents, chemistry, biochemistry and medicinal applications of different herbal essential oils and insights into the pharmacologically valuable plant proteases.

Chapter 1, written by Singh, illustrates the necessity for novel antivirals, which is further supported by the recent pandemic scenario. The effect of various Indian spices on the immune system, as well as how well they might be capable of fighting viral infections, is covered in detail in the second section.

Chapter 2, written by Zambelli, describes the chemical, functional and nutritional properties of Coriander.

In **Chapter 3**, Tiwari *et al.*, have compiled the traditional uses, chemical components and pharmacological properties of *Sphaeranthus indicus* Linn to provide a ready reference for scientists working on this plant species.

Chapter 4, written by Kumar *et al.*, comprehensively describes the phytochemistry of Copaiba oleoresins, conventional uses, and pharmacological properties of various oils from *Copaifera* species.

Chapter 5, written by Gupta *et al.*, comprehensively describes the pharmacognostic profile of *Nardostachys jatamansi*, a potent medicinal herb. In this chapter, they have described the various aspects of this medicinal plant in detail.

Chapter 6, written by OYETAYO *et al.*, describes a medicinal herb, Bush Onion (*Afrotyrax lepidophyllus*), along with its nutritional and medicinal values.

In **Chapter 7**, Agrawal *et al.*, describe the various plant’s essential oils, the chemistry of EO, extraction methods, essential oils in aromatherapy, as well as medicinal functional properties of different EOs like Lavender Oil, Clove oil, Eucalyptus oil, Peppermint oil, Lemongrass oil, and Cinnamon oil.

In **Chapter 8**, Roy *et al.*, have comprehensively discussed the various plant proteases, highlighting their pharmaceutical, industrial and biotechnological aspects as well as the structure, properties and catalytic mechanisms.

Pankaj Kumar Chaurasia
P.G. Department of Chemistry
L.S. College, B.R. Ambedkar Bihar University
Muzaffarpur-842001, Bihar, India

Shashi Lata Bharati

Department of Chemistry
North Eastern Regional Institute of Science and Technology
(NERIST) Nirjuli-791109
Arunachal Pradesh, India

&

Sunita Singh

Department of Chemistry
Navyug Kanya Mahavidyalaya, University of Lucknow
Lucknow, Uttar Pradesh, India

List of Contributors

Anuradha Singh	Department of Chemistry, Sadanlal Sanwaldas Khanna Girls' Degree College, Prayagraj, India
Arjun Patra	Department of Pharmacy, Guru Ghasidas Vishwavidyalaya, Bilaspur (C.G.), India
Brijesh Kumar	Department of Pharmacology, Institute of Medical Sciences, Banaras Hindu University, India
Folake Lucy Oyetayo	Department of Biochemistry, Ekiti State University, Ado-Ekiti, Nigeria
Hema Chaudhary	School of Medical & Allied Sciences, K.R. Mangalam University, Gurugram, India
Komal Rao	School of Medical & Allied Sciences, K.R. Mangalam University, Gurugram, India
Mona Piplani	Maharaja Agrasen School of Pharmacy, Maharaja Agrasen University, Atal Shiksha Kunj, Himachal Pradesh, India
Manmohan Singhal	Faculty of Pharmacy, DIT University, Dehradun, Uttarakhand, India
Mohit Agrawal	School of Medical & Allied Sciences, K.R. Mangalam University, Gurugram, India
Nathaniel Olawale Oluji	Department of Biochemistry, Ekiti State University, Ado-Ekiti, Nigeria
Nilasish Pal	Department of Chemistry, Seth Anandram Jaipuria College, Kolkata, India
Nirmalendu Das	Department of Botany, Barasat Govt. College, Barasat, Kolkata, India
Neetika Naudiyal	Department of Biochemistry, CBSH, GBPUAT, Pantnagar, Uttarakhand, India
Pratima Raypa	Department of Biochemistry, CBSH, GBPUAT, Pantnagar, Uttarakhand, India
Palak Gupta	Darjeeling Tea Research and Development centre, Tea Board, Kurseong (WB), Kurseong, India
Pankaj Bhateja	Maharaja Agrasen School of Pharmacy, Maharaja Agrasen University, Atal Shiksha Kunj, Himachal Pradesh, India
Rafael A. Zambelli	Food Engineering Department, Federal University of Ceará, Fortaleza, Brazil
Rajesh Kumar	Maharaja Agrasen School of Pharmacy, Maharaja Agrasen University, Atal Shiksha Kunj, Himachal Pradesh, India
Ruchika Garg	Maharaja Agrasen School of Pharmacy, Maharaja Agrasen University, Atal Shiksha Kunj, Himachal Pradesh, India
Shreyansi Das	Department of Botany, University of Calcutta, West Bengal, India
Sweta Arora	Plant Microbe Interaction and Molecular Biology Laboratory, Division of Biotechnology, CSIR Institute of Himalayan Bioresource Technology, Palampur, HP, India
Tina Roy	Plant Microbe Interaction and Molecular Biology Laboratory, Division of Biotechnology, CSIR-Institute of Himalayan Bioresource Technology, Palampur, HP, India

Vaishali Tiwari	Department of Pharmacy, Guru Ghasidas Vishwavidyalaya, Bilaspur (C.G.), India
Vandana A. Kumar	Department of Biochemistry , CBSH, GBPUAT, Pantnagar, Uttarakhand, India
Victor Olusegun Oyetayo	Department of Microbiology, Federal University of Technology, Akure, Nigeria
Varsha Yadav	Department of Pharmacy, Guru Ghasidas Vishwavidyalaya, Bilaspur (C.G.), India
Yogendra Singh	Maharaja Agrasen School of Pharmacy, Maharaja Agrasen University, Atal Shiksha Kunj, Himachal Pradesh, India

CHAPTER 1

Role of Indian Spices in the Management of Emerging Viral Infections

Anuradha Singh^{1,*}

¹ *Department of Chemistry, Sadanlal Sanwaldas Khanna Girls' Degree College, Prayagraj, India*

Abstract: The viral pandemic, with both emerging and re-emerging characteristics, has had a significant impact on human lives worldwide. This pandemic has resulted in substantial morbidity and mortality due to its rapid and widespread transmission among the general population. Unfortunately, the lack of effective antiviral treatments has further complicated the situation, making it challenging to control and manage the disease effectively. The ideal method for safeguarding against viral infections is to be immune. India's indigenous systems of medicine, with its rich array of phyto-therapies and natural remedies, have attracted the global community's interest. As a result, people all around the world are turning to and rekindling their interest in immune-boosting traditional remedies. Spices, with their significant therapeutic potential and excellent safety profiles, are currently being extensively studied as herbal drugs and immune-enhancing home remedies against infectious diseases. The first part of this chapter discusses the need for new antiviral drugs, especially in light of the recent pandemic. The second part reviews the extensive scientific evidence on the immune-boosting effects of many Indian spices, as well as their potential to fight viral infections. It also discusses their safety and toxicity profiles.

Keywords: Antiviral potential, Immune boosters, Safety profiles, Traditional remedies.

INTRODUCTION

From time immemorial, infectious diseases, particularly viral infections, have tormented humanity and civilization [1 - 4]. The virus's replication motifs and transmission modes are the prime roots of viral pandemics [5, 6]. However, substantial progress in the medical field against emerging viral infections has been made in the past few decades, particularly in the 1990s and 2000s. This progress has been driven by a number of factors, including advances in understanding the biology of viruses and their replication, the development of new technologies for

* **Corresponding author Anuradha Singh:** Department of Chemistry, Sadanlal Sanwaldas Khanna Girls' Degree College, Prayagraj, India; E-mail: anuradha_au@rediffmail.com

drug discovery and vaccine development, and increased investment in global health research and development [7 - 10].

The discovery of new treatment targets is not always followed by the formulation and authorization of new chemotherapeutics. Antivirals are not an exception when it comes to treating viral infections. Between 2017 and 2022, the FDA (Food and Drug Administration) granted approval to a mere seven new antiviral drugs, as highlighted in Table 1. The emerging and re-emerging viral outbreaks, as shown in Fig. (1), have propelled the scientific community to develop effective and sophisticated remedies to address the escalating demands imposed by these deadly diseases [11, 12].




Fig. (1). Emerging and re-emerging viral diseases.

Table 1. New antivirals approved by the FDA during the last five years (<https://www.fda.gov/drugs>).

Drug (Common Name)	Month and Year of Approval	Mechanism of Action	Treatment
Remdesivir (Veklury)	December 2020	Inhibition of RNA-dependent RNA polymerase enzyme of SARS-CoV-2	Treatment of COVID-19
Molnupiravir (Lagevrio)	November 2021	Oral antiviral drug that works by introducing errors into the RNA of SARS-CoV-2, which prevents the virus from replicating.	Treatment of asymptomatic or mildly symptomatic COVID-19 in high-risk adults

(Table 1) cont....

Drug (Common Name)	Month and Year of Approval	Mechanism of Action	Treatment
Paxlovid (PF-07321332)	December 2021	It is an oral antiviral drug that works by inhibiting the SARS-CoV-2 protease enzyme.	Managing the symptoms of mild-to-moderate COVID-19 in adults who are at high risk of severe disease
Evusheld (Tixagevimab and cilgavimab)	October 2021	This treatment works by using antibodies to target and neutralize, the SARS-CoV-2 spike protein, crucial for its attachment to human cells.	For the prevention of COVID-19 in adults and adolescents 12 years of age and older who are at high risk of severe disease.
Telaprevir (Invirase)	January 2022	It is an oral antiviral drug that works by inhibiting the protease enzyme of the hepatitis C virus.	For the treatment of chronic hepatitis C in adults who are co-infected with HIV.

Viral Life Cycle and Possible Drug Interventions

As parasites, contagious microscopic viruses encroach on living cells incorporated with host DNA. The virus takes control of the cellular machinery, initiating the production of numerous virion particles. As the entire life cycle of the virus unfolds within the host cell, discovering an effective treatment becomes exceedingly challenging. Different strategies have been developed by the scientific community to combat viral infections, and the main antiviral approaches currently under consideration may be classified into two types:

- (i) The approaches that directly target the viruses and
- (ii) The indirect approach involves targeting viruses by modulating the immune system of the host through various strategies, aiming to enhance the innate immune response or reduce inflammation triggered by viral infections [13].

The virus-directed methods require a thorough understanding of the virus's chemical nature, especially its interaction with the host cell. In the case of retroviruses, the exceedingly fast multiplication rate combined with the absence of exo-nuclease proofreading allows them to generate drug resistance mutations at an exceptionally fast rate. Viruses also exploit host machinery to not only replicate in host cells but also to escape the host immune system. The prospective targets in the viral life phases for the development of potential antivirals have been shown in Fig. (2).

CHAPTER 2

Chemical, Functional, and Nutritional Properties of Coriander (*Coriandrum Sativum*)

Rafael A. Zambelli^{1,*}

¹ Food Engineering Department, Federal University of Ceará, Fortaleza, Brazil

Abstract: Coriander is a plant common to several countries. Its use is associated with traditional medicine due to its functional and therapeutic properties. However, more and more coriander has been used in various sectors, such as food, medicine, pharmaceuticals, and cosmetics. Given the importance of this plant, this chapter aims to present the main chemical, functional and nutritional characteristics in addition to the main applications of coriander. The different parts of coriander have several bioactive components, such as monoterpenes, alkanes, aldehydes, alcohols, pigments, flavonoids, phenolic compounds, and saturated and unsaturated fatty acids. Because of its composition, coriander has numerous functional and nutritional properties such as antioxidant, antimicrobial, antibiofilm, antidiabetic, anxiolytic, and antidepressant activity, among others. Applications range from studying the impact on microorganisms to improving animal diets and extract preparations. Therefore, it can be said that coriander is a plant with multiple applications, and its nutritional potential should be increasingly explored to promote health benefits.

Keywords: Antimicrobial, Essential oil, Fatty acid, Monoterpenes, Phenolic compound.

INTRODUCTION

Several plants have aroused the interest of researchers because of their phytotherapeutic, medicinal and functional properties. In this context, an example is coriander. Because of the emerging concept of functional foods, it plays a role not just in nourishing the energy needed for human daily activities, but in cultivating healthy habits from a diet rich in bioactive components.

Coriander (*Coriandrum sativum* L.) is classified as a member of the *Umbelliferae* or *Apiaceae* family [1]. It is popularly known as a kind of condiment used by different cultures in cooking and is considered an edible plant [2]. Its origins are from the Mediterranean coast of Europe and Central Asia. It is especially popular

* Corresponding author Rafael A. Zambelli: Food Engineering Department, Federal University of Ceará, Fortaleza, Brazil; E-mail: zambelli@ufc.br

in Mexico, South America, and China. Coriander production has been increasing since 1994 due to increased consumer demand, especially in Asia, which is responsible for 71% of world production [3]. It is also grown naturally in North Africa [4].

In addition, there are reports of crops in Europe, especially in the Nordic countries (Finland, Denmark, Norway, and Sweden), Germany, Hungary, Belgium, and Poland [5]. The use of this plant has been recorded since before 1550 BC, being one of the oldest spice crops of humanity [6].

The top producers in 2022 were India (\$47.2M), Russia (\$29.0M), Italy (\$27.9M), (\$16.2M), (\$13.4M), Iran (\$10.8M), Ukraine (\$5.9M) and Canada (\$4.7M), according to a study [7].

Botanically, coriander is a slender, tender, glabrous, branched, annual plant and is considered a perennial herb that grows up to 50 cm in height. Its harvest period is between 2 to 3 months after sowing, and it is plucked by the roots [8].

This species has functional properties and nutraceutical and bioactive compounds of biotechnological interest [9]. In some cultures, coriander is planted with the aim of being a raw material for medicinal, culinary, and aromatic purposes. Its economic importance is valued due to its use as a flavoring agent in foods, cosmetics, and medicines [10]. Its leaves have an irritating and pungent aroma, and its seeds exude a sweet and spicy aroma [11]. The dried seeds are widely used as a spice in the Mediterranean region and India [12].

Dried fruits and leaves are used to provide aroma and flavor to food. Fruit is often used as a component of curry powder and is important in Indian cuisine. In turn, the leaves are often appreciated in Thai and Vietnamese cuisine [13]. In Brazil, especially in the northeast region, it is used as a seasoning in soups, broths, and beans. According to a study [14], the essential oils obtained from coriander fruits are used in food, alcohol, and cosmetic industries. Its leaves are rich in various food elements such as proteins, fats, fibers, vitamins, carbohydrates, water, vitamins C, calcium, phosphorus, iron, thiamine, riboflavin, and oxalic acid [15].

Research involving this plant focuses on the leaf, seed, or essential oil due to its attractive value in medicine and therapeutic potential; however, other parts of coriander may also have industrial potential.

Coriander is also highly valued for its functional and biological properties. Several studies have shown that parts of the plant may have antifungal, antiulcer, antimutagenic, anticancer, antiprotozoal, antidiabetic, anticonvulsant, antifertility, anxiolytic, diuretic, anthelmintic, sedation and hypnotic, hepatoprotective, and

cholesterol-lowering potential. It also has appreciable amounts of vitamins A, B, E and K, in addition to micro and macroelements and essential oils [16 - 19].

Coriander leaves have good amounts of vitamin C, carotenoids, dietary fiber, and polyphenols, as well as various amounts of volatile components that give coriander its specific fragrance. In turn, the seeds have high antioxidant activity, and this is attributed to the high contents of tocopherols, phospholipids, and carotenoids [20].

Chemical Properties

Green coriander contains 84% moisture, reduced content of saturated fats and cholesterol and high level of thiamine, zinc and dietary fiber [21], in addition to natural pigments such as chlorophyll-a, which gives the characteristic color of coriander leaves [22].

Coriander seeds have up to 1% essential oil and are composed of oxygenated monoterpenes, monoterpene hydrocarbons, long-chain alcohols, and fatty acids [23, 24]. A research points out that the monoterpenes present in coriander are linalool, camphor, geraniol and limolene.

The different parts of this plant contain monoterpenes, limpnene, α -pinene, γ -terpinene, p-cymene, citronellol, borneol, camphor, coriandrin, cis-geraniol, dihydrocoriandrin, coriandronsA-E, flavonoids, and essential oils [25]. Significant amounts of 2-Carene, p-mentha-1,3,8-triene, β -terpinyl-acetate, eucalyptol, β -myrcene, β -ocimene, trans-linalool oxide, terpinolene, borneol, terpinene-4-ol, α -terpineol, lavandulyl and geranyl acetate were found in coriander seeds [26].

In turn, compounds like decanal, (Z)-2-decenal, Undecanal, Dodecanal, 2-dodecenal, 2-tridecenal and 2-undecenal were found in the aerial parts of coriander; however, these components were not detected in the seeds [27]. These components are classified as aldehydes and can be formed from the oxidation of alcohols in the plant, which have antibacterial activity [28].

Many volatile components were found in coriander, including alcohols, aldehydes, hydrocarbons, and esters, with the class of aldehydes having the highest amount (878 $\mu\text{g}\cdot\text{g}^{-1}$) [29]. In turn, the presence of 46 aromatic components was detected in coriander essential oil, and the obtained coriander essential oil by the hydrodistillation method and gas chromatography analysis coupled with mass spectrometry identified 24 components, which were mostly aldehydes and alcohols [30].

CHAPTER 3

Phytochemical and Pharmacological Potential of *Sphaeranthus Indicus* Linn

Vaishali Tiwari¹, Varsha Yadav¹ and Arjun Patra^{1,*}

¹ Department of Pharmacy, Guru Ghasidas Vishwavidyalaya, Bilaspur (C.G.), India

Abstract: *Sphaeranthus indicus* Linn (Asteraceae) is commonly known as Munditika, Mundi, Shravana, Bhikshu, Tapodhana, Mahashravani, Shravanahva, and Shravanashirshaka. It is generally found in wet places of the lowlands and also grows as a weed in rice fields. Herein, the traditional uses, chemical components and pharmacological properties are compiled to provide a ready reference for scientists working on this plant species. The whole plant and various morphological parts are extensively used in Indian medicine to treat a variety of diseases. The plant has astringent, stomachic, reviving, pectoral, demulcent, and calming properties. Various secondary metabolites, such as eudesmanolides, sesquiterpenoids, sesquiterpene lactones, sesquiterpene acids, flavone glycosides, flavonoid C-glycosides, isoflavone glycoside, sterols, sterol glycoside, alkaloids, peptide alkaloids, amino acids, and sugars have been found in the plant. Numerous monoterpene hydrocarbons, oxygenated monoterpenes, oxygenated sesquiterpenes, and sesquiterpene hydrocarbons were found in the essential oils from the plant. The entire plant and various morphological parts have been reported to have ovicidal, anthelmintic, antifeedant, antimicrobial, antiviral, macrofilaricidal, larvicidal, analgesic, antipyretic, hepatoprotective, antitussive, wound healing, bronchodilatory, mast cell stabilising, anxiolytic, neuroleptic, immunomodulatory, anti-diabetic, antihyperlipidemic, antioxidant, central nervous system depressant, anti-arthritis, nephroprotective, and anticonvulsant activities. Acute toxicity of extracts from different plant parts revealed the safety of the drug.

Keywords: Phytochemistry, Pharmacological activity, *Sphaeranthus indicus*.

INTRODUCTION

Herbal medicine is the oldest kind of healthcare ever used by humans. Herbs have been used by all societies throughout history. Nowadays, herbal medicine is becoming more popular in both developed and developing countries. As improvements in clinical research, analysis, and quality control show how beneficial herbal medicine is in the treatment and prevention of disease, it is

* Corresponding author Arjun Patra: Department of Pharmacy, Guru Ghasidas Vishwavidyalaya, Bilaspur (C.G.), India; E-mail: drarjunpatra22@gmail.com

becoming ever more popular. Because they have fewer side effects, are much more easily accessible, are less expensive, *etc.*, herbal medicines are replacing synthetic ones to a greater extent frequently. Numerous drugs have entered the market as a result of research on ethnopharmacology and conventional therapies. Ayurveda, one of the Indian systems of medicine, employs a number of medicinal plants in the treatment of various disease conditions. The effectiveness of Ayurvedic medicine requires the establishment and validation of evidence. One of these medicinal plants that has been investigated for its phytopharmacological significance is *Sphaeranthus indicus* Linn.




Fig. (1). *Sphaeranthus indicus* **A)** Fresh whole plant of *S. indicus*, **B)** Fresh flower of *S. indicus*, **C)** Dried flower of *S. indicus*, **D)** Dried leaves of *S. indicus*.

The Hindi name of *S. indicus* is Gorakhmundi. It is a tall, multi-branched, fragrant plant that may grow up to 50 feet in the hills and is frequently found in India's plains (Fig. 1). The various morphological parts of the plant have been utilized in different forms against an array of health conditions. The plant is also rich in a variety of secondary metabolites having pharmacological significance. The information related to traditional uses, phytochemical investigation and

pharmacological studies on *S. indicus* was collected from different scientific databases such as Scopus, Pubmed, and Google Scholar and compiled here as a ready reference for future researchers to explore the plant from different angles.

PHYTOCHEMISTRY AND BIOACTIVE COMPOUNDS OF *SPHAERANTHUS INDICUS*

The alkaloid sphaeranthus with chemical formula $C_{13}H_{19}NO_5$ and melting point 166-168 °C has been isolated from chloroform extract of the entire *S. indicus* plant [1]. Additionally, they collected the volatile oil from *S. indicus* fresh flower by steam distillation and established its physicochemical constants. From the entire plant of *S. indicus*, three novel eudesmanoids (Fig. 2), 11,13-dihydro-3,7-dihydroxy-4,5-epoxy-6, 7-eudesmanolide (A), 11,13-dihydro-7-acetoxy-3-hydroxy-6 (B) and 7 eudesm-4-enolide and 3-keto- β -eudesmol (C) have been isolated from the whole plant of *S. indicus* [2].




Fig. (2). Novel eudesmanoids in *S. indicus*.

An immune-stimulating novel sesquiterpene glycoside called sphaeranthanolide has been isolated from the flowers of *S. indicus* and the structure was determined by using 2D-NMR and other spectroscopic methods (Fig. 3) [3]. Three novel eudesmanolides, 11,13-dihydro-3-,7-dihydroxyfrullanolide, 11,13-dihydro-7,13-dihydroxyfrullanolide and 11,13-dihydro-7-hydroxy-13-methoxyfrullanolide have been identified in the flowers of *S. indicus* [4]. A flavonoid C-glycoside called 5-hydroxy-7-methoxy-6-C-glycosylflavone has been isolated from the aerial parts of *S. indicus*, and spectroscopic techniques were employed to determine its structure [5]. From the entire *S. indicus* plant, the well-known 7-hydroxyeudesmanolide, two sesquiterpenoids (cryptomeridiol and 4-epicryptomeridiol), and 7a-hydroxyeudesmanolides have been identified, and structures of the novel compounds were determined by X-ray crystallography [6].

CHAPTER 4

The Chemistry and Pharmacological Study of *Copaifera* spp. Oils

Rajesh Kumar^{1,*}, Yogendra Singh¹, Mona Piplani¹, Pankaj Bhateja¹, Ruchika Garg¹ and Brijesh Kumar^{2,*}

¹ Maharaja Agrasen School of Pharmacy, Maharaja Agrasen University, Atal Shiksha Kunj, Himachal Pradesh, India

² Department of Pharmacology, Institute of Medical Sciences, Banaras Hindu University, India

Abstract: Since ancient times, *Copaifera* tree oleoresin has been widely utilised for conventional medicine, and it is currently a well-liked remedy for a variety of illnesses. The vast majority of the chemical makeup that makes up copaiba resins is frequently composed of sesquiterpene hydrocarbons such as germacrene D, copaene, caryophyllene, elemene and humulene. Kaurenoic acid, alepterolic acid, copalic acid and polyalthic acid are a few of the physiologically useful diterpene compounds present in oleoresin. Due to its many pharmacological qualities and extensive usage, oleoresin is one of the most significant regenerative natural remedies and folk medicines. In addition to being used as a contraceptive, oil or plant bark decoction is also used to treat inflammation, bronchitis, syphilis, and cough. Wound healing is improved when oil is administered topically to the skin. During massages, it is administered to the head to manage convulsions, pain, and paralysis. It is also suggested to apply oil-soaked cotton to tumours, ulcers, or hives. Dog bites and other infected wounds are treated with the bark's decoction, which is also used to treat rheumatism. The oil is utilised to create a patch that is mechanically applied to wounds and some ulcers to treat them. Caryophyllene has exhibited its activity in a number of pharmacological models, including cannabinoid receptors, making it one of the most significant phytoconstituents in copaiba oils at the present time. The goal of this chapter is to look into the phytochemistry of copaiba oleoresins, conventional uses, and the pharmacological properties of various oils from *Copaifera* species.

Keywords: β -caryophyllene, Bark decoction, Bronchitis, Convulsions, Diterpene acids, Inflammation, Sesquiterpenes, Oleoresin, Paralysis, Tumours.

* Corresponding authors Rajesh Kumar & Brijesh Kumar: Maharaja Agrasen School of Pharmacy, Maharaja Agrasen University, Atal Shiksha Kunj, Himachal Pradesh, India & Department of Pharmacology, Institute of Medical Sciences, Banaras Hindu University, India; E-mails: rajdhiman60@gmail.com, asthwal@rediffmail.com

INTRODUCTION

According to the taxonomical classification of the genus *Copaifera*, it belongs to the subfamily *Caesalpinoideae* and family *Fabaceae*. The copaiba trees make up the bulk of the *Copaifera* genus' tree species [1]. The allusions to the indigenous terms copava and copahu, which refer to a plant's secretions and the oil found within it, respectively, are most likely to be applicable to the origin of basic and common copaba [2].

In 1638, Marcgraf and Piso used the term “Copaiba” to describe the genus for the first time without identifying the particular species. Some scientists mentioned the plant species *Copaiva officinalis* in their work *Enumeratio Systematica Plantarum* in 1760. In his subsequent investigations, *Species Plantarum*, which was printed in 1764, Carl von Linnaeus accomplished a more comprehensive examination of the plant and specifically named it *Copaifera officinalis* (Jacq.) L as a particular type [3]. There are around 70 *Copaifera* species, with South and Central America possessing the largest abundance. In addition, a total of four species may be found in the African continent, and one subspecies can be found in Borneo, an island in the Pacific. Brazil has the most *Copaifera* biodiversity, with 26 species and 8 variations [4].

In light of the various botanical synonyms associated with certain species, more thorough, environmentally friendly, taxonomical, genomic, metabolomic, and genetic studies on the genus are necessary, as well as a revaluation of the species classification [5]. The botanical features of the leaves, such as the presence or lack of notches on the leaflets, the amount and shape of the leaflets, and floral attributes, such as the lengths of the petals and anthers and the condition of the stamens, are the primary distinguishing features among various *Copaifera* species [6]. To basically outline these plants, they can be recognised botanically through their petiolate and terminal leaves. These trees can live for a maximum of 400 years, grow slowly, exhibit shrubby or arboreal inclinations, reach heights of up to 40 m, and have peak diameters of up to 4 m [7].

Intercellular glandular lines are arranged in zones of peripheral axial parenchyma across their cylindrical shafts. The parenchyma cells of the tube's parenchyma produce a substance called the secretory layers' lumen and develop schizogenously. The species' leaves are alternating, pinnate, and 2 to 12 paired. Generally, clear points and glands at the base of the bordering vein may occasionally be present [8]. The species is frequently deciduous and possesses thin, interpetiolar stipules as well. The interior hirsute sepal containing a tetra calice of the monoclamid, small, many, sessile flowers creates short tubes. Small bracts provide protection for the flower buds, and the spikes of flowers are

alternating panicles. The gynoecium features a sessile in nature receptacle with two elongated eggs, a filiform fashion, and a cylindrical and papillary stigma as opposed to the androecium, with its ten loose stamens that are smooth fillets, and a rectangular, rimose anther [9].

The fruits are laterally compressed, dehiscent, bivalved, and monospermic. The seed is pendulous, oblong-globular, and densely coated in white or yellow arils, and the endosperm is absent. The blooms are minor, apetalous, hermaphrodite, and grouped in axillary panicles. The rectangular seed of the fruits is encircled by a swarm of vivid arils [10].

Although the *Copaifera* genus has undergone extensive taxonomic examination, certain species remain difficult to discern, mostly because of their delicate floral morphology and the lack of reproductive systems in the samples studied. The primary obstacle to botanical identifying the collection with respect to Amazonian species is the paucity of field data and specimen pictures [11]. The development of programmes, strategies, and plans for the long-term preservation and conservation of commercially important species has also been hampered by these taxonomic issues. These issues have also impeded the advancement of chemical and medical research, as well as the industrial and practical uses of resin oils and the use of wood [12].

Significant sources of fatty acids and timber are the aforementioned trees. The liquid known as oleoresin varies in colour (from yellow to light brown) and viscosity. Its non-volatile component, which is mostly made up of acid compounds, and its dynamic component, which is a combination of sesquiterpenes, are both solid forms and polymers [13].

Numerous traditional remedies employ this oleoresin. The molecules with the kaurane, clerodane, and labdane skeleton types can be found in all oleoresins despite the fact that the species and individuals within the same species exhibit substantial variability in the chemical profile of the oleoresin. It is notable that copalic acid, a labdane-type diterpene, is present in the majority of industrial samples of the drug [14].

The major components of the volatile fraction are thought to be sesquiterpenes that have two forms, oxygenated and non-oxygenated, such as caryophyllene, copaene, humulene, and caryophyllene oxide [11]. Researchers in this area have shown the most interest in the *Copaifera langsdorffii* L. After the leaves' hydroalcoholic extract was subjected to phytochemical analysis, flavonoid heterosides such as kaempferol-3-o-alpha-L-rhamnopyranoside (afzelin), galloylquinic and methylated galloylquinic acids, quercetin-3-o-alpha-L-rhamnopyranoside (quercitrin), and others were discovered [15].

CHAPTER 5

Pharmacognostic Profile of a Potent Medicinal Herb: *Nardostachys Jatamansi*

Palak Gupta^{1,*}, Neetika Naudiyal², Pratima Raypa² and Vandana A. Kumar²

¹ Darjeeling Tea Research and Development centre, Tea Board, Kurseong (WB), Kurseong, India

² Department of Biochemistry, CBSH, GBPUAT, Pantnagar, Uttarakhand, India

Abstract: *Nardostachys jatamansi* DC, a small, erect, perennial, rhizomatous herb growing on steep, moist, rocky, undisturbed grassy slopes, is a popular species of aromatic and medicinal plant. The Sanskrit word “Jata,” which denotes a matted and uncombed lock of hair, is similar to its rhizome, hence the name “*Jatamansi*.” The plant is found between 3000 and 5200 m above sea level in the alpine Himalayas, which include parts of Uttarakhand, Himachal Pradesh, Arunachal Pradesh, and Sikkim in India, as well as numerous regions of Nepal, Tibet, China, and Bhutan.

N. jatamansi has been used in ayurveda, ethnomedicine and alternative medicine for a very long time, going all the way back to the Vedic era (1000–800 B.C.). Its rhizomes were used as a bitter tonic to cure epilepsy, stimulant, antispasmodic and to treat hysteria in the traditional Ayurvedic medicinal system. It has a wide range of therapeutic benefits, including anti-inflammatory, antidepressant-like action, anticonvulsant, hypotensive, anti-asthmatic, anti-estrogenic, cardioprotective, neuroprotective, antidiabetic, antifungal, antibacterial, and anticancer characteristics. Additionally, it functions as a sedative, tranquillizer, hypolipidemic, anti-lipid peroxidative, antioxidant, and anti-oxidant. In addition, it is used to treat a variety of neurological conditions, including insomnia, excitation, epilepsy, neurosis, Alzheimer's disease, and issues with learning and memory. As per various phytochemical studies, esters, phenolic chemicals, and terpenic ketone are the major components present. Due to the frequently observed variations in the types of metabolites in the plant, tremendous medicinal applications can be precisely explored using the database of antioxidant, antimicrobial, anticancer, antidiabetic and anti-inflammatory activities, making it a potential candidate for treating oxidative stress and other diseases produced during pathogenic conditions. For the prevention and treatment of a variety of pathological illnesses, as well as the side effects associated with them, the use of plants like *N. jatamansi* may thereby lessen reliance on synthetic pharmaceuticals.

* Corresponding author Palak Gupta: Darjeeling Tea Research and Development centre, Tea Board, Kurseong (WB), Kurseong, India; E-mail: palakbiochem801@gmail.com

Keywords: Antioxidant, Antimicrobial, Antidiabetic, Jata, Oxidative stress, Perennial.

INTRODUCTION

People have relied on nature for healthcare for millennia, including plants, insects, animals, and fungi. Communities have found several plant species with varied medical applications over time and have maintained significant ethnobotanical knowledge to improve the quality of their lives. Despite the fact that current science has discovered plants and plant extracts that can treat and cure ailments, it may have been challenging for prehistoric humans to identify and classify plants having components beneficial to health. According to the earliest known medical records between 5000 and 3000 BC, Sumerian clay tablets show that people were aware of ailments and thus used plants with medicinal properties to protect and restore health. People were aware of the medical plants, as evidenced by the medicinal herbs found on the preserved remains of Tzi, the Iceman, who was unintentionally murdered between 3400 and 3100 BC in the icy, high Alps. Plants probably flourished by producing disgusting, unpleasant chemical components that repulsed animals, as they may be susceptible to destruction owing to being used as fodder by animals and insects [1]. Humans may have developed the selective practice of eating particular plant parts as a result of their observations that various plant tissues, such as fruits, leaves, or roots of particular plants, improved their mood. From these foundations, lovely plant gardens have been created for the food and plant components that contribute to people's continued health.

The World Health Organization (WHO) defines “conventional medicine” as the totality of knowledge, skills, and practices derived from theories, beliefs, and stories that are precise to numerous cultures and used in the maintenance of health as well as through preventive and therapeutic measures for physical and mental illness. Because of its potential and effective applications and minimal side effects, herbal treatment for many ailments is currently gaining popularity. In the human community, medicinal and aromatic plants (MAPs) are a significant source of medications and are being used as a main component in the field of healthcare. MAPs are a large group of species with a variety of sources, traits, and functions that are a component of traditional medical systems that may be found in many local cultures all over the world. The usefulness of medicinal and aromatic plants depends on their accessibility and active therapeutic capability to manage particular illnesses. Each medicinal and aromatic plant contains a unique kind and quantity of chemical components that vary according to the plant species. Additionally, when plants get older, their concentration changes. According to Zhang's report 1996 [2], the significance of traditional medicinal herbs in the

prevention or control of some metabolic disorders like diabetes, heart disease, and specific types of malignancies has been amply demonstrated by research-based data. Since ancient times, herbal medicines have been utilized in treating a variety of illnesses throughout the world [3, 4]. It has been documented that medicinal plants can be used alone, in combination, or even as the main source of raw materials for other medications [5].

Higher plant species, which represent by far the highest use of the natural world in terms of species, have been used in traditional and modern medicine all over the world in amounts ranging from 50,000 to 70,000 species, with many more species important to the expanding market for plant-based cosmetics and other products. These species now play a crucial role in healthcare in many poor nations, delivering effective medicines for sizable segments of the population in places where other types of medication are either unavailable or unaffordable. According to estimates, 80% of the populations in Africa and Asia rely heavily on these plant-based medications for their healthcare requirements, and the WHO [6] predicted that in the ensuing decades, a similar percentage of the global population may do the same. The majority of the MAP trade takes place on a local level within countries, and the majority of these species are only used in traditional medicine [7 - 9]. Approximately, 3000 of these plant species have been traded globally. An estimated 1,77,000 metric tons of herbal medicine were traded in India each year that included 960 plant species [10]. Around 2500 medicinal plants from the Indian subcontinent are used locally for medical purposes and/or are traded, particularly in the pharmaceutical industries. Of these, 1748 species have their origins in the Indian Himalayan Region, a majority of which are found in the Uttarakhand region [11].

Nardostachys jatamansi is a medicinal and aromatic plant local to the Himalayan region that has an extended record of use in conventional Indian remedies and historic Greco-Arab remedies. Due to *N. jatamansi*'s medicinal and aromatic properties, 300 metric tons of raw materials (dry rhizome) are needed in India each year [10].

Botanical Classification

Contrary to what is stated in all the documented material about this plant, *N. jatamansi* actually belongs to the *Caprifoliaceae* family, not the Valerianaceae.

Kingdom: Plantae

Division: Mangnoliophyta

Class: Mangnoliopsida

CHAPTER 6

Potentials of Wild Edible Bush Onion (*Afrostryax Lepidophyllus*) Plant as a Source of Essential Food and Medicinal Constituents

Victor Olusegun Oyetayo^{1,*}, Folake Lucy Oyetayo² and Nathaniel Olawale Oluji²

¹ Department of Microbiology, Federal University of Technology, Akure, Nigeria

² Department of Biochemistry, Ekiti State University, Ado-Ekiti, Nigeria

Abstract: The Bush Onion (*Afrostryax lepidophyllus*) plant is an unconventional tropical West African plant characterized by a strong onion-like aroma. It is a member of the Huaceae family, and its bark is used as food sauces, condiments, spices, and flavorings due to its strong aroma. The seed is used trado-medicinally to treat venereal diseases and gastro-enteric diseases and as a laxative and mouthwash. It also has anti-cancer, antioxidant, estrogenic and insecticidal properties. These effects have been attributed to its content of phytochemical compounds such as polyphenols, flavonoids, glycosides, saponins, alkaloids, and anthocyanins. It also contains appreciable concentrations of certain important phytonutrients. Its potential as an affordable nutritive source of non-synthetic constituents can be exploited as a valuable contribution to food and medicine sources, especially in the developing world.

Keywords: Antioxidant, Bush onion, Estrogenic, Medicinal, Phytonutrients, Unconventional.

INTRODUCTION

The 'Bush Onion' plant (*Afrostryax lepidophyllus*) is a wild edible plant that grows in some parts of tropical Africa and is characterized by a strong onion/garlic-like aroma. The tree bark is used in food as sauces, condiments, spices, and flavorings. The seed is believed to possess anti-cancer activity and is repulsive to poisonous animals such as snakes and scorpions [1]. It is used as a febrifuge to treat venereal and gastro-enteric diseases and as an antidote for snake bites [2]. The root bark decoction is consumed as an anthelmintic and an enema to treat urinary infections [3], while the bark is used to treat convulsions in Ghana [4].

* Corresponding author Victor Olusegun Oyetayo: Department of Microbiology, Federal University of Technology, Akure, Nigeria; E-mail: ovonew67@gmail.com

The plant extract has been shown to have antioxidant activity, estrogenic properties and insecticidal effects [5], as shown on the cowpea weevil, *Callosobruchus maculatus* [6]. The essential oils extracted from the bush onion seed were reported to exhibit cytotoxic and antimicrobial properties [7].

The seed contains important food nutrients that can contribute significantly to the daily nutrient requirement of all age groups. It is an important potential flavoring agent in food processing with several industrial and commercial applications. As indigenous people globally depend on plants obtainable from the forest, the bush onion seed, though edible and underutilized, is a potential source of readily available natural phytoconstituents that can alleviate malnutrition and provide an array of substances of medicinal importance, which can be exploited as a medicinal food.



Plate. (1). *Afrostyrax lepidophyllus* plant. Source [8]:



Plate. (2). *Afrostyrax lepidophyllus* dried seeds. source [8]:

Botanical Description and Historical Cultivation

Afrostyrax lepidophyllus, commonly known as 'country onion' or 'bush onion plant', belongs to the family Huaceae [2]. Plate (1) shows the *Afrostyrax lepidophyllus* plant [8]. It is a non-timber forest product. The tree grows up to 20 m high in the green forests of the Democratic Republic of the Congo, Nigeria, Ghana, the Central African Republic, and Cameroon, where its natural existence is becoming threatened due to the conversion of natural forests into commercial plantations and in some cases mining activities. The tree, leaves and fruit have a strongly offensive smell of onion or garlic, and the bark slash is particularly strong. The oval-shaped fruit [8] has a pleasant aroma (Plate 2). The seeds ripen between July and August but appear as vulnerable on the International Union for Conservation of Nature's (IUCN) red list [9].

Chemical and Nutritional Composition of Bush Onion

The seed oils, as well as tree bark of *Afrostyrax lepidophyllus*, showed the presence of sulfur-containing compounds, mainly thio-ethers with 2,4,5,7-

CHAPTER 7

Essential Oil: Its Chemistry, Biochemistry and Application

Mohit Agrawal^{1,*}, Komal Rao¹, Manmohan Singhal² and Hema Chaudhary¹

¹ School of Medical & Allied Sciences, K.R. Mangalam University, Gurugram, India

² Faculty of Pharmacy, DIT University, Dehradun, Uttarakhand, India

Abstract: Essential oils are secondary metabolites that are produced by a variety of medicinal plants. These are mostly composed of a blend of terpenic hydrocarbons and oxygenated derivatives like ketones, alcohols, epoxides, aldehydes, and esters. These are insoluble in water but are soluble in ether, alcohol, and fixed oils. The majority of these volatile oils are liquid and colourless at room temperature. The majority of the components found in essential oils are from the terpene family and are concentrated in the cell secretions of specific plants. The chemical makeup and positioning of functional groups on molecules affect the mechanism of action of essential oils. When it comes to their antibacterial activity, essential oils' chemical makeup influences how they work. The content is about several essential oils, such as lavender oil, which has antibacterial and antifungal properties due to its primary ingredients, linalool and β -ocimene. Both anti-inflammatory and antibacterial properties can be found in eucalyptus oil, which contains the primary ingredients 1,8 cineole, cryptone, p-cymene, and pinene. Sulphur molecules found in garlic essential oils are effective in preventing cancer. In order to confirm the safety and effectiveness of essential oils, more clinical trials are obligatory. Essential oils possess the ability to be employed as defensive as well as therapeutic agents for a variety of ailments.

Keywords: Constituents, Essential oil, Efficacy, Metabolites, Terpene.

INTRODUCTION

Essential oils are metabolites (secondary metabolites) that are developed by a diversity of plants having medicinal value. These are mostly composed of a combination of terpenic hydrocarbons and compounds like ketones, alcohols, epoxides, aldehydes, and esters [1 - 4]. These are obtained by fermentation, expression, effleurage or extraction of different plants having medicinal value.

* Corresponding author Mohit Agrawal: School of Medical & Allied Sciences, K.R. Mangalam University, Gurugram, India; E-mail: mohitagrawalmohu@gmail.com


Aside from these procedures, hydro distillation and steam distillation are the most often employed procedures for producing essential oils [5, 6]. Approximately 3000 EOs have been discovered so far. These are primarily found in the flavedo part and cuticle oil glands or sacs that are present at various depths in fruit peels [7, 8]. Essential oils are derived from different parts of plants, including barks, seeds, flowers, leaves, and peels. They have antifungal, antiviral, antioxidant, and antibacterial properties and can be employed as safe alternatives to harmful chemical preservatives and substances that promote food preservation. Since ancient times, the consumption of essential oils for various dental as well as medical issues has been documented [9 - 12].

The compositions of EOs vary significantly. Even the chemical makeup of EOs isolated from plants belonging to the same species varies according to location. The constitution of the EOs also relies on how mature the plant is from which they are harvested, and their use is still widespread but restricted to the fields of cosmetics and perfumes. These are insoluble in water and can dissolve in ether, alcohol, and fixed oils [13]. At room temperature, the majority of these volatile oils are liquid and colourless. With the exception of a few instances, they smell distinctively and have a density that is lower than that of cinnamon, sassafras, and vetiver). They are very optically active and have a refractive index. Herbs contain volatile oils, which are the source of the various aromas that plants release. These are frequently employed in perfumery, cosmetics, and aromatherapy fields. The latter includes massage, inhalations, or baths employing these volatile oils as a therapeutic approach [14 - 17].


CHEMISTRY OF ESSENTIAL OILS (EOs)

Production of essential oils involves a variety of distinct structures; the quantity and make-up are highly varied. EOs are concentrated in the cytoplasm of specific secretions of plant cells, which are found in more than one organ plant [18, 19]. Essential oils are intricate combinations that include more than 300 distinct chemicals [20]. They are primarily composed of organic volatile chemicals with low molecular weights (under 300). They are found in part in the vapour state due to their high vapour pressure at atmospheric pressure and room temperature [21].


Essential oils are composed of hydrocarbon molecules, like other organic chemicals, and can be further divided into phenols, alcohol, terpenes, esters, aldehydes, and ketones, among other categories. Numerous other aromatic compounds are available from aldehydes, alcohols, and ketones, including fruity ((E)-nerolidol), herbal (-selinene), floral (Linalool), citrus (Limonene), and more as shown in Fig. (1) [22, 23]. Some structures related to the chemistry of essential oils are [24 - 28]:



Structure of Terpenoids



Structure of Geraniol



Structure of Chamomile

Fig. (1). Some structures related to the chemistry of essential oils [24 - 28].

Terpenoids

The main elements of essential oils produced by several species of plants and flowers are terpenes and terpenoids. The monoterpene and sesquiterpene

CHAPTER 8

The Plant Proteases-A Group of Potential Pharmacological, Biotechnological, and Industrially Important Enzymes

Tina Roy¹, Sweta Arora¹, Shreyansi Das², Nilasish Pal^{3,*} and Nirmalendu Das^{4,*}

¹ Plant Microbe Interaction and Molecular Biology Laboratory, Division of Biotechnology, CSIR-Institute of Himalayan Bioresource Technology, Palampur, HP, India

² Department of Botany, University of Calcutta, West Bengal, India

³ Department of Chemistry, Seth Anandram Jaipuria College, Kolkata, India

⁴ Department of Botany, Barasat Govt. College, Barasat, Kolkata, India

Abstract: Proteases play an active role in all living organisms. In plants, they exhibit essential roles in various biological processes throughout the life cycle by proteolysis, like cell growth and development, differentiation, seed germination, reproduction, senescence, development of stomata, chloroplast biogenesis, apoptosis, plant resistance, *etc.* Nowadays, it is a well-known group of enzymes for its industrial and therapeutic values. Plant proteases *viz.* papain (papaya), bromelain (pineapple), ficin (fig tree), and cucumisin (melon) are well-characterized proteases that have medicinal values for the treatment of upset stomach, diabetic thrombosis, wounds, tumor, asthma, angina, rheumatoid arthritis, bronchitis, sinusitis, *etc.* In addition, some common Indian spices are found to be good sources of natural proteases, such as *Withania coagulans* (vegetable rennet), *Carum copticum* (Ajwain), *Syzygium aromaticum* (Clove), *Cuminum cyminum* (Cumin), *Nigella sativa* (black cumin), *Cinnamomum verum* (Cinnamon), *Foeniculum vulgare* (common fennel), *Zingiber officinale* (ginger), *Cinnamomum tamala* (Indian Cassia), *Curcuma longa* (turmeric), *etc.* which have traditional usage as healing agents in multiple disorders. This chapter highlights the pharmaceutical, industrial and biotechnological aspects of various plant proteases as well as the structure, properties and catalytic mechanisms.

Keywords: Bromelain, Cucumisin, Enzymes, Peptidase, Proteinase, Proteolysis, Spices.

* Corresponding authors Nilasish Pal and Nirmalendu Das: Department of Chemistry, Seth Anandram Jaipuria College, Kolkata, India; and Department of Botany, Barasat Govt. College, Barasat, Kolkata, India; E-mails: nilasishpal@gmail.com and nirmalendus@yahoo.co.uk

INTRODUCTION

Proteases (E.C.3.4.21.14) are a set of enzymes that catalyze the breakdown of peptide linkages in proteins through hydrolysis and are generally referred to as proteolytic enzymes, peptidases or proteinases. Proteases are omnipresent in biological systems and are required for cellular metabolism [1]. Different groups of hydrolytic enzymes come under proteases, which are classified on the basis of their site of action, structure of the active site and specific reaction mechanisms [2]. Proteases play a major role in regulating cellular functions in all living organisms, including the human body, which incorporates cell differentiation, division and cell death. Also, they are biochemically involved in acquiring nutrition, growth, and adaptation. Intracellular proteases are involved in protein degradation processes using lysosomes and proteosomes [2]. Plants, animals and especially microorganisms are the foremost sources of proteases.

Proteases account for 65% of total enzymes, considered as enzymes with high industrial demand for their role in food, detergent, leather processing, and chemical and pharmaceutical applications [3, 4]. Proteases are enzymes used in several therapeutic applications, including diagnosis of diseases like treating wounds, cancer, and digestive disorders. In the food industry, it has been used for different processes such as brewing, clotting of milk and meat tenderizing. Plant proteases can be obtained with cost-effective production and are safe as they do not carry pathogenic infectious agents [5]. They have an estimated market value of 3 billion US\$, with a compound annual growth rate of 6.1% by the year 2024 [6].

Proteases are found in almost all living beings starting from simple prokaryotes to higher eukaryotes. The majority of proteases are obtained from microbial sources, but newer technologies are being developed to utilize plant sources as well. In plants, the most abundant among them is cysteine protease, while aspartic protease is the rarest. Proteases have been obtained and studied extensively from the latex of a number of dicot plant families, such as Asteraceae, Apocynaceae, Asclepiadaceae, Caricaceae, Euphorbiaceae and Moraceae [7].

Plant-derived proteases are also now being explored as an alternative for calf rennet which is a complex of different proteases and is frequently used in cheese making [8]. In recent years, the demand for cheese has increased, whereas calf rennet production has decreased, which has led to an increase in the price of cheese as well as paved the way for finding new and appropriate rennet substitutes [9]. Different animal, microbial and plant sources for proteases were studied for the same. The development of plant proteases for the production of cheeses is suitable for vegetarians. Different plant parts such as stem, fruit, and leaves of

Withania coagulans, *Carica papaya*, *Ananas comosus*, *Ficus carica* and *Actinidia deliciosa* are being used to produce conventional plant proteases such as bromelain, ficin, papain, actinidin, *etc.* and are involved in meat tenderization, dairy industry, baking industry, animal feed, biomedicine, alcohol production and many other industrial processes [1]. Besides these plant sources, research on new plant proteases with higher industrial efficiency through *in vitro* culture or genetic engineering is still ongoing.

CLASSIFICATION OF PROTEASES

Proteases are broadly classified into two categories: “endopeptidases” and “exopeptidases” according to their position of action, *i.e.*, where they cleave the peptide bond. The endopeptidases are further categorized by their catalytic types. Exopeptidases are also divided into two classes according to whether they act at the N- or C-terminus (Fig. 1).




Fig. (1). Schematic illustration of different Protease families and multidimensional uses (Blue-Exopeptidases; Red- Endopeptidases).

SUBJECT INDEX**A**

- Acid 20, 25, 26, 28, 29, 51, 52, 58, 86, 87, 88, 89, 90, 91, 92, 98, 99, 127, 129, 137, 151, 160, 180, 185, 186, 190, 200
- arachidic 26
- arachidonic 137
- ascorbic 51
- aspartic 186
- benzoic 90
- butyric 200
- catic 86, 88, 90
- chlorogenic 25, 129
- chrolechinic 86, 90, 91
- elaidic 87, 92
- gallic 25, 52, 90, 92, 129
- glutamic 180, 185, 186, 190
- hypochlorous 127
- linoleic 26, 58
- lipoic 127
- oxalic 20
- palmitic 26, 28, 29
- pentanoic 86, 89, 90
- petroselinic 26, 29
- protocatechuic 129
- sinapic 26
- Activity 19, 32, 51, 54, 68, 69, 70, 72, 95, 97, 101, 102, 128, 135, 165
- antibiotic 97
- anticancer 72, 95, 102, 165
- antidepressant 19
- antiproliferative 135
- anxiolytic 54
- enzymatic 32
- fungicidal 101
- immunomodulatory 69, 72
- iron chelating 51
- iron chelation 128
- nephroprotective 70, 72
- neuropharmacological 68
- Agents 4, 84, 107, 151, 164, 168, 179
- antidepressant 168
- antiemetic 151
- antiviral 4
- immunoregulatory 164
- microbiological 84
- Alanine 66, 67
- aminotransferase 66, 67
- transaminase 66
- Allergic reactions 137, 161, 168
- Alzheimer's disease 117, 120, 136
- Aminopeptidases 178
- Analgesic 65
- activity 65
- effectiveness 65
- Angiotensin-converting enzyme (ACE) 179, 186
- Anti-diabetic 61, 136
- effectiveness 61
- effects 61, 136
- Anti-inflammatory 9, 12, 31, 62, 63, 72, 95, 99, 100, 117, 137, 152, 164, 196
- activity 62, 63, 72, 100, 117, 137, 196
- effects 9, 31, 62, 99
- properties 12, 95, 100, 137, 152, 164
- Anti-lipidemic action 100
- Anti-nutrient composition 151
- Anti-oxidant effects 51
- Anti-termite effects 96
- Anti-tumor actions 97
- Antibacterial 50, 51, 52, 55, 56, 57, 58, 95, 97, 101, 102, 132, 133, 157, 158, 162, 163, 164, 165
- activity 50, 51, 55, 56, 57, 58, 95, 101, 132, 133, 157, 164, 165
- effect 97, 164
- properties 52, 56, 57, 58, 97, 102, 157, 158, 162, 163, 164
- Antibiotic resistance 132
- Anticancer effects 102
- Anticonvulsant effects 70
- Antidepressant effects 11

Subject Index

Antidiabetic 19, 20, 29, 35, 60, 61, 72, 117, 118, 120, 136, 137
 action 61
 activity 60, 72
 agent 35
Antiepileptic properties 30
Antifeedant effects 64
Antifungal 7, 55, 56, 57, 58, 95, 96, 152, 157, 164, 165, 168
 action 96
 activities 55, 56, 58, 95, 152, 164, 165
 agent 152
 properties 7, 57, 157, 168
Antihyperglycemic action 136
Antileishmanial activity 103
Antimicrobial 27, 29, 33, 34, 35, 54, 55, 56, 57, 58, 72, 95, 132, 133, 148, 165
 activity 29, 33, 34, 54, 55, 56, 57, 58, 72, 132, 133
 agent 35
 properties 27, 58, 95, 148, 165
Antimutagenic effects 29, 95
Antinociceptive effect 65
Antioxidant 8, 23, 27, 28, 51, 52, 53, 72, 127, 128, 137, 148, 150, 152, 164
 activities 23, 28, 51, 52, 53, 72, 128, 137, 148, 152, 164
 agent 52
 enzymes 8, 127
 properties 27, 51, 52, 53, 127, 150, 152
Antiproliferative effects 95, 135
Antispasmodic effects 120
Antitumor 195
 activity 195
 effects 195
Antiviral 4, 5, 10
 action 10
 drugs 4
 treatments 5
Apoptosis 96, 97, 134, 175, 178, 195
Aromatherapy oils 161
Aspartate 66, 67
 aminotransferase 66, 67
 transaminase 66
Aspartic 176, 179, 180, 189, 190, 199
 peptidases 180
 proteases 176, 179, 180, 190, 199
 proteinases 189
Aspartyl glucosaminidase 187

The Chemistry Inside Spices & Herbs, Vol. 4 213

Atherosclerosis 62
Ayurvedic medicine 45

B

Bacterial adhesion 101
Blood vessel reticulocytes 102
Bromelain monotherapy 196
Bronchitis 79, 175, 195, 196, 199
Bronchodilatory activity 64
Bronchospasm 64, 69
 acute 64
 histamine-induced 64

C

Cancer 8, 30, 60, 96, 127, 133, 134, 136, 151, 153, 164, 176
 breast 60
 lung 96
Carcinogenesis 134
Carcinoma 95, 96, 135
 breast 135
 malignant 96
 renal 95
Chromatography 22, 30, 50, 55, 98, 105
 gas 22, 50, 98, 105
 thin-layer 55, 105
Chymotrypsin 178, 186
Cinnamomum tamala 175
Conditions 117, 127, 200
 chronic 127
 neurological 117
 plant growth 200
COVID-19 pandemic 6, 7
Cysteine 176, 179, 180, 186, 188, 191, 195
 peptidases 191
 proteases 176, 179, 180, 186, 188, 195
Cytokines 9, 63
 anti-inflammatory 9
 pro-inflammatory 63
Cytotoxicity activity 96

D

Dental caries 97, 151
Dietary fiber 21
Diffusion assay 55
Diseases 1, 3, 5, 6, 7, 8, 13, 44, 72, 119, 120,

127, 132, 133, 153, 164, 176, 179, 182,
183, 194, 195, 196, 197
autoimmune 194
cardiovascular 127, 196
celiac 182
heart 119, 153
infectious 1, 6, 132, 164
mental 120
neurodegenerative 8
skin 72, 197
Disorders 5, 7, 8, 83, 119, 120, 127, 131, 152,
195
degenerative 152
hypersensitivity 195
metabolic 5, 8, 119
nervous system 120
neurodegenerative 127
neurological 8
respiratory 83
viral infectious 7

E

Enzyme(s) 10, 11, 12, 28, 52, 136, 175, 176,
178, 179, 180, 184, 185, 186, 192, 193,
194, 198, 199, 200, 201
angiotensin-converting 186
inhibition 52
inflammatory cell 136
polymerase 11
polyphenol oxidase 28
proteolytic 176, 185, 192
Epidermidis 55, 56, 57, 58, 132, 133
Erythrocytes 102, 107
micronucleated polychromatic 107
Erythromycin 57
Essential oil(s) (EOs) 20, 21, 50, 130, 152,
157,
158, 159, 160, 161, 162, 165, 166, 167,
168
and aromatic water 161
in aromatherapy 166
producing 158
Esterolytic activity 182
Ethnopharmacology 45

F

Fish protein hydrolysis 181

Flight mass spectrometry 66
Fluorescence microscopy 60
Food 34, 126, 162, 168, 176, 178, 181
additives 126, 162
digestion 178
industry 34, 168, 176, 181
Foot-shock induced aggression (FSIA) 54
FRAP assay 51
Fuels 85, 184
fossil 85
Fumigation bioassay 103

G

Galloylquinic acid derivatives 105
Gas chromatography 21, 104, 105, 129
analysis 21
approach 104
mass spectrometry 129
technique 105
Gastrointestinal adenocarcinoma 96
Genetic manipulations 201
Genotoxicity 107, 127
Gentamycin 56
Glucose absorption 61
non-insulin-dependent 61
Glutamate 186, 191
protease 186
residue 191
Glutamic proteases 180, 190, 191
Glutathione transferase 53

H

Hepatic glucose production 136
Hepatitis, chronic 3
Hepatocellular degeneration 67
Hepatoprotective 67
agent 67
effects 67
mechanisms 67
High-performance liquid chromatography 129
Human respiratory syncytial virus (HRSV) 10
Hydrogen peroxide assay 69
Hydrolysis reaction 180

I

Immune 1, 7, 10, 178

Subject Index

boosters 1, 7
 response 10, 178
Immunity 6, 8, 69, 168
 boosting 8
 humoral 69
Immunomodulatory 7, 69
 action 69
 effects 7, 69
Industries 20, 35, 36, 84, 177, 181, 197, 200, 201
 baking 177, 181, 197
 cosmetic 20
 wine 181, 197
Infections 6, 10, 13, 32, 82, 132, 151, 194, 196, 199
 bacterial 32, 82
 intestinal 194
 urinary tract 151
Inflammation 3, 8, 9, 11, 12, 31, 63, 79, 96, 127, 136, 137
 formalin-induced chronic 63
Insulin 61, 136, 166
 plasma 6
 production 166
 resistance 136
 sensitivity 136
Iron chelation 51

L

Lemongrass essential oil (LEO) 165
Lipid profiles 60, 61, 136
Lipoxygenase 137
Liver 30, 66, 67
 damage 66, 67
 disorders 30
Low-density lipoproteins (LDL) 60, 61, 62, 66

M

Mass spectrometry (MS) 21, 22, 31, 125, 129
Metabolic profiling 128
Metabolism 61, 107
 lipid 61
 mitochondrial energy 107
Metabolite profiling 128, 129
Metal-ligand enzyme complexes 131
Metalloproteinases 185, 191

The Chemistry Inside Spices & Herbs, Vol. 4 215

Metalloproteinases 179, 183, 185, 191, 192
Metals, toxic 32
Metastasis 153, 195
Methicillin-resistant *S. aureus* (MRSA) 58, 133
Methods, ion-exchange chromatography 98
Milk protein 179
Mutagenesis effects 107
Mutagenic and cytotoxic activity 102

N

Neuroblastoma 134
Neuropeptide processing 178
Neuropharmacological effects 68
Neuroprotective effects 68

O

Oils 29, 162, 164, 167
 clove 162, 164
 crude 29
 lemon grass 167
Oleic acid 26, 29
Osteoarthritis 12, 199

P

Pain regulation 12
Pathway, mitochondrial apoptotic 195
Photographic industry 85
Plant(s) 28, 36, 118, 119, 124, 126, 134, 175, 176, 177, 178, 180, 195, 196, 197, 198, 201
 aromatic 118, 119
 environment 126
 expression systems 198
 hormones 28
 metabolites, secondary 134
 nutraceutical 36
 proteases 175, 176, 177, 178, 180, 195, 197, 198, 201
 proteolytic enzymes 196
 tissue culture technique 124
Pneumonia 56, 57, 58, 132, 133
Pro-inflammatory leukotrienes 137
Processes 31, 128, 132, 136, 160, 161, 176, 177, 178, 179, 180, 185, 191, 195, 198, 199, 200

- enzymatic 161
- fermentation 185
- hormonal regulation 185
- industrial 177
- inflammatory 31, 136
- Products 68, 165, 167
 - cosmetic 165, 167
 - fluorescence 68
- Properties 7, 8, 9, 12, 29, 31, 34, 35, 61, 64, 69, 101, 103, 153, 165, 166, 167, 168, 175, 178
 - anesthetic 29
 - antihyperglycaemic 61
 - antileishmanial 103
 - antimutagenic 166
 - antiproliferative 153
 - bronchodilatory 64
 - carcinogenic 168
 - chelating 35
 - immune-protective 7
 - immunity booster 7
 - immunomodulatory 9, 12, 69
 - immunostimulatory 31
- Protease(s) 3, 175, 176, 177, 178, 179, 180, 181, 183, 184, 185, 186, 193, 194, 196, 197, 198, 201
 - activity 185
 - enzymes 3, 179, 194, 196
 - microbial 185
 - plant-derived 176, 201
- Protein(s) 20, 34, 55, 124, 127, 128, 135, 150, 152, 176, 178, 180, 185, 186, 192, 197, 200
 - amyloid precursor 180
 - autoproteolytic 192
 - collagen 197
 - hydrolysis 197
 - metabolism 135
 - nucleophiles 185, 186
 - tumor suppressor 135
- Proteinases 175, 176, 193
- Proteolysis 175, 186, 187, 191, 195, 197
- Proteolytic activity 185

R

- Reactive oxygen species (ROS) 8, 22, 27, 127, 136, 153
- Reducing power assays 69
- Regulators, metabolic 130

- Respiratory distress 10
- Response 6, 12, 66, 68, 126, 128, 166, 195
 - humoral immune 12
 - immunological 6
 - reducing inflammatory 195
 - robust analgesic 66
- Rheumatoid arthritis (RA) 63, 175, 194, 199
- RNA-dependent RNA polymerase enzyme 2
- Rosemary oil 167

S

- Seed germination 175, 180
- Serotonin pathways 100
- Serum 60, 61, 62, 66, 67, 71, 196
 - alanine aminotransferase 67
 - creatinine 62
 - glutamate pyruvate transaminase 62
 - glutamic oxaloacetic transaminase 62
 - insulin 61
- Silicon supplementation 23
- Skin 167, 168
 - complexion 167
 - greasy 167
 - oily 168
- Stress 30, 31, 68, 117, 118, 126, 127, 152, 166, 167
 - chronic 30
 - induced neurotoxicity 68
 - oxidative 31, 117, 118, 127, 152
- System 3, 4, 7, 9, 12, 96, 167
 - immune 3, 4, 7, 9, 12, 167
 - lymphatic 96

T

- Technique 46, 49, 64, 130
 - hemocytometer 64
 - plant protection 130
 - spectroscopic 46, 49
- Therapeutic 8, 83, 106, 157, 169, 194
 - agents 157, 169, 194
 - effects 8, 83
 - plants 106
- Thermolysin 185, 186, 192
- Total 52, 60, 61, 62, 66, 69, 128
 - cholesterol (TC) 60, 61, 62, 66
 - phenolic content (TPC) 52, 69, 128

Traditional medicinal systems 5
Transcription factor 9
Tumors 31, 134, 135, 137, 175
 antiproliferative 134
 necrosis factor 31, 137
Tyrosinase 52

V

Vaginal cream 106
Virus(s) 2, 3, 6, 10, 11, 12
 dengue 12
 herpes simplex 11
 influenza 10, 12


W

Waste management 185
Water stress 22
Wound 31, 44, 53, 54, 79, 178, 182, 183, 194,
 195
 constriction 53
 healing 31, 44, 54, 79, 178, 182, 183, 194,
 195
Wound healing 53, 54, 72, 100, 101
 activity 53, 54, 72, 100
 effect 54
 procedures 101



Pankaj Kumar Chaurasia

Dr. Pankaj Kumar Chaurasia did Ph.D. in chemistry and is an assistant professor in the PG Department of Chemistry, L.S. College, B.R. Ambedkar Bihar University, Muzaffarpur. He has experience of more than 6 years in teaching at post graduate (PG) and undergraduate (UG) levels. He qualified the 'National Eligibility Test' in 2009 as CSIR-JRF (NET) and was awarded with SRF-NET in 2012. He also worked as guest faculty in the Department of Chemistry, Central University of Allahabad, Prayagraj. He was also awarded with 'Research Associateship' by CSIR-New Delhi (India) and worked as CSIR-RA in Motilal Nehru NIT, Allahabad/Prayagraj (India). He has more than 73 publications. He is also guiding four Ph.D. research scholars. He has worked in different areas of chemistry and has research expertise in biological chemistry, biotechnology, microbiology, enzymology, organometallic chemistry and environmental chemistry.



Shashi Lata Bharati

Dr. Shashi Lata Bharati, did Ph.D. in chemistry and is currently working as assistant professor in the Department of Chemistry, North Eastern Regional Institute of Science and Technology, Nirjuli, Arunachal Pradesh (India). She was awarded 'UGC-DSA fellowship' for the meritorious students during her Ph.D. program. She was also awarded with a UGC-post doctoral fellowship for women by UGC New Delhi and worked as post-doctoral fellow in the Department of Chemistry of DDU Gorakhpur University, Gorakhpur (India). She has published more than 50 publications in national and international journals/publishers of repute. She has guided many M.Sc. project students and one Ph.D. research scholar. She has expertise in the field of inorganic chemistry, organometallic chemistry, biological chemistry and enzymology.



Sunita Singh

Dr. Sunita Singh is an assistant professor of chemistry in the Navyug Kanya Mahavidyalaya since 2019. She pursued her Ph.D. in the year 2015 in chemistry. She was awarded with JRF and SRF from the University Grant Commission. She also worked as research assistant (2010-2011) on a CST sponsored project. She has published 25 research articles in journals of national and international repute. She has authored and co authored 8 book chapters with national and international publishing houses. She has two edited books with international publishers. She is also on the reviewer board of different journals. She delivered many invited talks and attended several conferences/webinars of national and international repute. She is also an active member of Association of Chemistry Teachers (India), International Clinical aromatherapy network and Global Harmonization Initiative. She has expertise in chemistry, phytochemistry, essential oil, oleoresin etc.